

Possible Evidence for a Chiral Particle B_0^χ , Scalar Meson in the B Meson System

Daiki ITO, Muneyuki ISHIDA*, Shin ISHIDA, Toshihiko KOMADA**, Tomohito
MAEDA and Ichiro YAMAUCHI***

*Research Institute of Quantum Science, College of Science and Technology
Nihon University, Tokyo 101-0062, Japan*

**Department of Physics, Tokyo Institute of Technology, Tokyo 152-8551, Japan*

***Department of Engineering Science, Junior College Funabashi Campus
Nihon University, Funabashi 274-8501, Japan*

**** Tokyo Metropolitan College of Technology, Tokyo 140-0011, Japan*

In the covariant level-classification scheme of hadrons proposed recently, the existence of “chiral particles”, scalar and axial-vector mesons as the partners of the ground-state pseudo-scalar and vector mesons, respectively, were predicted in the D meson and B meson systems, realizing a linear representation of chiral symmetry. In this work we reanalyze the $B^0 \pi$ mass spectra through Z^0 decays to show a possibility for the scalar B_0^χ .

§1. Introduction

The level classification of hadrons has been done rather successfully on the basis of the non-relativistic quark model(NRQM) for these 4 decades. However, this non-relativistic scheme seems now to be confronted with a serious difficulty: Recently existence^{1), 2)} of the light-mass iso-scalar scalar meson to be identified with σ ^{3), 4)}, the chiral partner of the π meson as a Nambu-Goldstone boson, has been accepted widely⁵⁾. Successively possible existence of the $\kappa(900)$ meson⁶⁾, iso-spinor scalar meson, has been pointed out. Then we are able to identify naturally the σ -meson nonet⁷⁾ with the members $\{\sigma(600), \kappa(900), a_0(980), f_0(980)\}$.

The mass values ($\leq 1\text{GeV}$) of this scalar nonet are in the region of $q\bar{q}$ ground states in NRQM, while there are no seats for scalars in this model. This has aroused recently a hot controversy on what the quark configuration of σ nonet is.

On the other hand we have proposed a new level-classification scheme^{8), 9)}, which has a relativistically covariant framework and in conformity with the approximate chiral symmetry. In this new classification scheme the “chiral states”, which are out of the framework of the NRQM, are expected to exist in the lower mass region, and the σ -nonet is naturally assigned as the $(q\bar{q})$ relativistic S -wave chiral state. In the heavy-light quark meson system it is expected that the approximate chiral symmetry on the light quark is valid, and the new level-classification scheme anticipates existence of the chiral states¹⁰⁾, scalar mesons and axial-vector mesons, in addition to the conventional states, pseudo-scalar and vector mesons, in the ground states.

In the previous work¹¹⁾ we have shown a possible evidence for existence of a chiral particle D_1^χ , axial-vector in the D meson system. The purpose of this paper is to investigate the possibility for existence of another chiral particle B_0^χ , scalar in

the B meson system, by reanalyzing some experimental data reported previously.

§2. Experimental Data and Project for Reanalysis

(*Experimental data to be analyzed*)

As the experimental data to be reanalyzed we choose the mass spectra of $B\pi$ system obtained through Z^0 boson decay

$$e^+ + e^- \rightarrow Z^0 \rightarrow B\pi + \dots, \quad (2.1)$$

by L3 collaboration¹²⁾ with some statistical accuracy. We also apply the data by ALEPH collaboration¹³⁾ with low statistical accuracy for supplementary analysis. The original data are inclusive and the relevant exclusive mass spectra of $B\pi$ system are obtained by subtracting from the original ones the backgrounds of the form

$$B. G. = P_1(\Delta M)^{P_3} \exp \left[P_4(\Delta M) + P_5(\Delta M)^2 + P_6(\Delta M)^3 \right] \quad (2.2)$$

$$\Delta M \equiv M_{B\pi} - P_2,$$

with fitting parameters P_i ($i = 1, 2, \dots, 6$). We shall apply the same formula to Eq. (2.2) for the backgrounds, taking into account the possible effects of intermediate B_0^χ production.

(*Relevant intermediate resonances*)

The conventional two $(b\bar{q}) - P$ wave mesons may contribute directly, as intermediate states, to the final $B\pi$ system with the respective angular momenta $l = 0, 2$ as

$$\text{Direct Reson. Process} \quad B_0^* \rightarrow B + \pi \quad (S - \text{wave}), \quad (2.3)$$

$$B_2^* \rightarrow B + \pi \quad (D - \text{wave}), \quad (2.4)$$

where B_0^* and B_2^* have, respectively, $^{j_q}L_J = ^{1/2}P_0$ and $^{3/2}P_2$ ($\mathbf{j}_q = \mathbf{S}_q + \mathbf{L}$ is the total angular momentum of the light quark).

In this work, a possible direct contribution from, in addition to the above conventional resonances, the chiral scalar meson B_0^χ is taken into account as

$$\text{Direct Reson. Process} \quad B_0^\chi \rightarrow B + \pi \quad (S - \text{wave}), \quad (2.5)$$

Concerning a possible candidate for the chiral particle B_0^χ , some excess structure over the backgrounds around the mass $m \sim 5550$ MeV, which is common to both the data of L3 and ALEPH, is identified as being due to production of B_0^χ .

In the relevant experiment the low energy γ was unable to be observed. Accordingly we must take into account the background process from the intermediate resonances, decaying into $B^* + \pi$ (successively B^* decays into B and missing γ).

$$\text{Backgd. Reson. Process} \quad B_1^* \rightarrow B^* + \pi \quad (S - \text{wave}), \quad (2.6)$$

$$B_1 \rightarrow B^* + \pi \quad (D - \text{wave}), \quad (2.7)$$

$$B_2^* \rightarrow B^* + \pi \quad (D - \text{wave}), \quad (2.8)$$

$$B^* \rightarrow B + \gamma. \quad (2.9)$$

(Formulas for analysis)

We shall apply the VMW method in our relevant case, where the absolute amplitude squared is given by

$$|M(s)|^2 = \{|r_1 \Delta_{B_0^\chi}(s) + r_2 e^{i\theta} \Delta_{B_0^*}(s)|^2 + |r_3 \Delta_{B_2^*}(s)|^2\} \\ + \{|r_4 \Delta_{B_1^*}(s)|^2 + |r_5 \Delta_{B_2^*}(s)|^2 + |r_6 \Delta_{B_1}(s)|^2\} \quad (2.10)$$

$$\Delta_i(s) \equiv \frac{-m_i \Gamma_i}{s - m_i^2 + im_i \Gamma_i} \quad (i \text{ denoting respective resonances}) . \quad (2.11)$$

Here the first (second) term represents the contributions from the direct (background) resonance processes Eqs. (2.3-2.5) (Eqs. (2.6-2.8)), the $m_i(\Gamma_i)$ are the mass(width) of relevant resonances, and r_i represent their production strength. In Eq. (2.10) a possible interference effect between the two direct decay processes of the B_0^* (Eq. (2.3)) and of the B_0^χ (Eq. (2.5)) are taken into account, while no interference effects among any background process Eqs. (2.6-2.8) are expected.

(Project for applying χ^2 -analysis)

We shall perform the two kinds of χ^2 -fitting of the mass spectra, the one with B_0^χ and the other without B_0^χ , and compare with their results. Our main interest for reanalysis in this work is to search for a possibility of existence of B_0^χ . Accordingly we shall follow to the projects of original analysis, modifying them only in the places to be related with this existence, and our projects are:

- i) Backgrounds; in the case without B_0^χ their parameters, P_1 through P_6 , are determined so as to reproduce the original background curve given by L3, while in the case with B_0^χ the original curve is revised in the region close to threshold where the possible effects of B_0^χ may be included.
- ii) mass and width of resonances; their values of m_i and Γ_i appeared in Formulas (2.10) are restricted in the regions given in Table I. In this table $M_{B_1^*}$ and $M_{B_2^*}$ are treated as free parameters and the restrictions $M_{B_0^*} = M_{B_1^*} - 12\text{MeV}$ and $M_{B_1} = M_{B_2^*} - 12\text{MeV}$ are set up. Also $\Gamma_{B_1^*}$ and $\Gamma_{B_2^*}$ are free parameters and the restrictions $\Gamma_{B_0^*} = \Gamma_{B_1^*}$ and $\Gamma_{B_1} = \Gamma_{B_2^*}$ are set up. It is also taken into account the fact that the effective mass of relevant resonances in the background processes is shift down by $\Delta M_\gamma = m_{B^*} - m_B = 46\text{MeV}$. The upper and lower limits of the regions on the free parameters $m_{B_1^*}$, $m_{B_2^*}$, $\Gamma_{B_1^*}$ and $\Gamma_{B_2^*}$ are determined in reference to their original values of L3 also shown in Table I.

Table I. Resiticted regions of masses and widths (in MeV)

Direct Res. Process			Background Res. Process		
B_0^χ	m	5530–5570	B_1^*	m	5590–5640
	Γ	10–40			(L3; $5670 \pm 10 \pm 13$)
B_0^*	m	$m(B_1^*) - 12 + 46$		Γ	20–100
	Γ	$\Gamma(B_1^*)$			(L3; $70 \pm 21 \pm 25$)
B_2^*	m	5730–5780	B_1	m	$m(B_2^*) - 12 - 46$
		(L3; $5768 \pm 5 \pm 6$)		Γ	$\Gamma(B_2^*)$
	Γ	0–60	B_2^*	m	$m(B_2^*) - 46$
		(L3; $24 \pm 19 \pm 24$)		Γ	$\Gamma(B_2^*)$

- iii) Energy resolution of experiments; its effects are taken into account through

the relation

$$\Gamma_{R_i} = \sqrt{\Gamma_{i,\text{orig.}}^2 + \Gamma_{i,\text{resol.}}^2} , \quad (2.12)$$

where $\Gamma_{i,\text{resol.}}$ is the energy resolution σ of L3, Γ_{R_i} is the value for our formula, and $\Gamma_{i,\text{orig.}}$ is the true width of the resonance R_i .

iv) Production couplings r_i ; following L3, the production ratios among resonances are assumed to be proportional to the $(2J_i + 1)$ (J_i being spin of R_i). Then r_i are obtained from Γ_i through the relation

$$r_i^2 \Gamma_i \propto (2J_i + 1) . \quad (2.13)$$

§3. Results of Analysis

The results of our analysis in both cases, with B_0^χ and without B_0^χ , are shown in Fig. 1. The fitted curves in comparison with the data are given, respectively, as follows: In the case with B_0^χ [without B_0^χ] the results on exclusive L3 spectra and ALEPH ones are shown in Fig. (a), (c) and (e) [(b), (d) and (f)] .

The values of mass and width of direct resonances, B_0^χ , B_0^* and B_2^* , obtained through our analysis of (a) L3(exclusive) are given in Table II. In Table III the

Table II. Fitted values of mass and width (in MeV)

	B_0^χ	B_0^*	B_2^*
m	5534	5635	5743
Γ	19.7	40.3	73.1

values of χ^2 and of $\tilde{\chi}^2 (\equiv \chi^2 / \text{No. of data points} - \text{No. of param.})$ of the analysis of L3(exclusive) are given in both the cases with and without B_0^χ . Through the

Table III. Values of χ^2 and $\tilde{\chi}^2$

	with B_0^χ	without B_0^χ
χ^2	18.5	22.4
$\tilde{\chi}^2$	$\frac{18.5}{34-9} = 0.74$	$\frac{22.4}{34-5} = 0.77$

inspection of fitted curves in Fig. 1, we may conclude that our results give some evidence for existence of B_0^χ , although there is no significant improvement on the reduced χ^2 .

§4. Concluding Remarks

(*Summary of this work*) We have made reanalysis of the $(B\pi)$ mass spectra obtained through the process Eq. (2.1), by applying the formulas (2.10) and (2.11). As the intermediate direct resonance we have taken into account the B_0^χ , a new chiral scalar meson outside of the conventional level-classification scheme. As a result we have obtained a possible evidence of B_0^χ with $m = 5530$ MeV and $\Gamma = 19.7$ MeV. The obtained values of reduced χ square, $\tilde{\chi}^2 = 18.5/(34 - 9) = 0.74$ for the case with B_0^χ and $22.4/(34 - 5) = 0.77$ for the case without B_0^χ , respectively. However, the




(a) L3(exclusive) with B_0^χ (b) L3(exclusive) without B_0^χ (c) L3(inclusive) with B_0^χ (d) L3(inclusive) without B_0^χ (e) ALEPH with B_0^χ (f) ALEPH without B_0^χ

Fig. 1. Fitted curves in the case with (without) B_0^χ and experimental data of L3(exclusive), L3(inclusive) and ALEPH are shown, respectively, in Fig. (a), (c) and (e) ((b), (d) and (f)). The contributions from respective resonances are also shown.

statistical accuracy of the relevant data is very poor, and the more accurate data are required in order to get a definite conclusion.

(*Universal property of chiral particles*) In the previous works the universal relations^{10), 14), 15)} of chiral particles through the D - and B -meson systems are predicted from the chiral symmetry on the light-quark and the heavy quark symmetry: The mass splittings between the respective chiral partners are equal, and the decay widths for the one-pion emission of the chiral particles, scalar and axial vector mesons are also equal.

We can check some of these universal relations experimentally, using the results of the analyses given in this letter and the previous one¹¹⁾, as follows:

$$\begin{aligned} \Delta m_D (\equiv m(D_1^\chi) - m(D^*)) &= \Delta m_B (\equiv m(B_0^\chi) - m(B)) \\ 302(2312 - 2010) \text{ MeV} & \quad 238(= 5535 - 5297) \text{ MeV}. \end{aligned} \quad (4.1)$$

$$\begin{aligned} \Gamma(D_1^\chi \rightarrow D^* \pi) &= \Gamma(B_0^\chi \rightarrow B \pi) \\ 23 \text{ MeV} & \quad 20 \text{ MeV}. \end{aligned} \quad (4.2)$$

From the above we may conclude that the theoretical predictions on the universal properties of chiral particles are consistent with the present experiment.

The authors should like to express their gratitude to Professors, K. Takamatsu and T. Tsuru, for their useful discussions and encouragement.

References

- [1] R. Delbourgo and M. D. Scadron, Phys. Rev. Lett. **48** (1982), 765.
- [2] T. Hatsuda and T. Kunihiro, Prog. Theor. Phys. **74** (1985), 765; Phys. Rep. **247** (1994), 221.
- [3] N. N. Achasov and G. N. Schestakov, Phys. Rev. D **49** (1994), 5779.
R. Kaminski, L. Lesniak and J. -P. Maillet, Phys. Rev. D **50** (1994), 3145.
N. A. Tornqvist, Z. Phys. C **68** (1995), 647.
N. A. Tornqvist and M. Roos, Phys. Rev. Lett. **76** (1996), 1575.
M. Harada, F. Sannino and J. Schechter, Phys. Rev. D **54** (1996), 1991.
- [4] S. Ishida, M. Ishida, H. Takahashi, T. Ishida, K. Takamatsu and T. Tsuru, Prog. Theor. Phys. **95** (1996), 745; **98** (1997), 1005.
- [5] Proceedings on *Possible Existence of σ -Meson and Its Implications to Hadron Physics*, YITP, Kyoto, 2000 (“ σ -Meson 2000”), Nihon University Preprint, NUP-B-2000-1; So-ryushiron Kenkyu (Kyoto) **102** (2001), E1; KEK-proceedings/2000-4.
- [6] S. Ishida, M. Ishida, T. Ishida, K. Takamatsu and T. Tsuru, Prog. Theor. Phys. **98** (1997), 159.
- [7] M. Ishida, Prog. Theor. Phys. **101** (1999), 661.
- [8] S. Ishida, M. Ishida and T. Maeda, Prog. Theor. Phys. **104** (2000), 785.
- [9] S. Ishida and M. Ishida, Phys. Lett. **B 539** (2002), 249.
- [10] M. Ishida and S. Ishida, Prog. Theor. Phys. **106** (2001), 373; hep-ph/0110360.
- [11] D. Ito, M. Ishida, S. Ishida, T. Komada, H. Tonooka and K. Yamada, to be published.
- [12] L3 collaboration, S. Goldfarb et al., Phys. Lett. **B465** (1999), 323.
- [13] ALEPH collaboration, S. Monteil et al., in Proc. of “Workshop on Hadron Spectroscopy”: Frascati Physics Series XV (1999) p.305.
- [14] W. A. Bardeen and C. T. Hill, Phys. Rev. D **49** (1994), 409.
- [15] D. Ebert, T. Feldmann, R. Friedrich and H. Reinhardt, Nucl. Phys. B **434** (1995), 619.